



PLATFORM TENNIS NEWS

Official Publication of the American Platform Tennis Association

THE SOUND OF A GUNSHOT

by Stephanie Frey

Gregg Brents has been playing professional platform tennis since 1974. Like most players on the circuit, he has had his share of scrapes and bruises and minor injuries. However, he was unprepared for what happened to him on March 27, 1992.

Brents and his partner, Andy Kinney, were at double match point in the third set of the first round of the President's Cup competition in Huntington, Long Island. Brents went back for an overhead and suddenly heard what sounded like a gunshot from a small-caliber pistol. Brents went down for the count, His partner missed the point.

Brents had heard this sickening sound five years earlier when playing with Carter Reul. He knew immediately what had happened.

His wife Jan drove him straight to the office of **Dr. Jacob Rozbruch**, chief of orthopaedic surgery at Beth Israel North Medical Center. Although Brents was not a patient of **Dr. Rozbruch's**, Jan was. **Dr. Rozbruch is a specialist in sports-related injuries.**

Dr. Rozbruch confirmed his worst fears. Brents had completely ruptured his Achilles tendon. The good news was that he would make a 100-percent comeback. The bad news it would not be for six to nine months. This meant Brents would not see the beginning of the platform tennis season in October and would be lucky to be back on the court by January.

Achilles Tendon Ruptures

Achilles tendon ruptures are not uncommon among 40 to 50-year-old recreational tennis players. (Brents is 41.) Sports such as tennis, football, or basketball, which require repeated stop-and-go movements, are much more likely to result in an Achilles tendon injury than long-distance running or skiing, for instance. While the average Achilles tendon rupture is not sports related, top professional athletes are not immune from the injury, e.g., NFL's Lawrence Taylor of the New York Giants and NBAs Dominique Wilkins of the Atlanta Hawks.

The Achilles tendon and gastroc-muscle complex are primarily responsible for quick acceleration and long stride lengths. In racket sports in particular, the muscle complex and tendon are continuously stretched and contracted in rapid succession. Constant repetition of this movement may, over time, cause degeneration of the tendon, possibly resulting in a rupture through the degenerated portion.

On the platform tennis circuit, torn Achilles tendons have been a recurring problem. Several top players - Brents' former partner Carter Reul, Phil Crane, and, most recently, Buddy Scudder - have suffered similar injuries.

According to **Dr. Rozbruch**, "Achilles tendon ruptures are not difficult to diagnose if you know what you are looking for." However, it is not uncommon for the injury to be misdiagnosed by non-orthopaedists.

Dr. Rozbruch recounts an instance when a patient came to him for a second opinion with an Achilles tendon injury that had been diagnosed at an hospital emergency room as an ankle sprain. Upon examination, **Dr. Rozbruch** realized the mistake and immediately treated the problem with surgery. Unfortunately, the patient had already been walking on the injury for two to three weeks prior to the discovery, thereby lengthening the recovery period.

Brents' former partner Carter Reul experienced a similar situation. The initial diagnosis for his injury was a slightly torn Achilles tendon. Based on this analysis, Ruel walked around for eight weeks thinking the injury would mend itself. During that time, significant scar tissue had built up. On the advice of a second doctor, he had it operated on, after which it took Reul almost a year to return to the game 100 percent.

Rozbruch counsels that injuries of this nature should be handled by an orthopaedic specialist not a general practitioner or emergency-room physician. "Not every emergency room physician is orthopaedically trained," states **Dr. Rozbruch**.

Dr. Rozbruch also cautions those who suffer from Achilles tendinitis. "While it is not difficult to diagnose the problem, the suggested treatment is not always accurate." The worst solution is to inject the tendon with cortisone. "Cortisone shots may reduce inflammation and pain but will ultimately weaken the tendon, making a full rupture more likely to occur," he advises.

"A full rupture in an athletic individual," **Rozbruch** believes, "merits surgery. A partial rupture in an athletic or non-athletic patient can be treated with casting. An inflamed Achilles' tendon should be treated with rest."

For Brents' torn Achilles tendon, **Dr. Rozbruch** recommended surgical repair followed by casting and aggressive physical therapy.

The operation took one and a half hours. The procedure involved surgically sewing Brents' tendon back together and augmenting it through localized grafting taken from the patient's own tendons. This added extra strength. Upon completion, Brents was fitted with a hip-to-toe cast to immobilize and protect the leg and encourage healing of the surgery.

One day later, Brents was discharged from the hospital. One week later, the full leg cast was removed and replaced with a below-the-knee cast, which he wore for another seven weeks.

Recovery & Rehab

According to **Dr. Rozbruch**, the prognosis for recovery is determined by the type of injury. “If it’s very close to the bone, the more difficult it is to get a good result,” states **Rozbruch**. “If it’s up near the muscle, the better the results.” In Brents’ case, the injury was located in mid-tendon.

In addition to the type of injury and surgical technique, individual fitness, both physiological and psychological, are key factors in determining the length of recovery.

Mitzi Gerard, co-owner of Advanced Physical Therapy and Sports Rehab who has a master’s of science in physical therapy, says that Brents’ extraordinarily rapid recovery was primarily due to the active role he took in his rehabilitation. “He was an eager and willing participant,” Gerard states.

Upon removal of the cast, Gerard started Brents off with a fairly aggressive physical-therapy program that was to last three months. The initial phase consisted of hydrotherapy in a whirlpool, high-volt electrical stimulation, heat, ultrasound, deep-connective-tissue massage to stretch out the fibers in the muscle and work the tendon, range of motion exercises and therapeutic activities to strengthen and increase circulation to the area:

Gerard’s main goal during this time was to get the foot, which had been held in a downward or plantar flexed position, back to a neutral position so Brents could walk normally again. This first stage of recovery lasted six to eight weeks.

For the final phase of rehab, Gerard put Brents on the Biodex isokinetic-resistance equipment to build power and strength. This, along with skill-specific start-and-stop exercises in the advanced stages of therapy, enabled Brents to get back on the court.

Brents’ recovery was exceptionally fast. With this therapeutic approach, the normal time period before a patient recovering from Achilles tendon surgery can return to an aggressive, competitive level of play is generally six months. Brents took three.

“I have a 10-inch zigzag scar down the back of my leg,” he says. “It looks vicious, but I have no pain and full mobility.”

Dr. Rozbruch had several reasons for using the zigzag incision in Brents’ particular case. First was to avoid possible scar contracture that would have impeded Brents’ recovery process, and secondly, to avoid risking the formation of a keloid or thickened scar.

“Zigzag incisions help eliminate the axial pull on the incision during the healing process by shifting the tension away from the scar to the surrounding skin,” he explains. “In this manner, the elasticity of the skin can be optimized without ripping the scar.” **Dr. Rozbruch** reserves this technique for athletes who require an accelerated recovery period and maximum stretch during play. He would not recommend it, for instance, to people who are concerned about its cosmetic aspects.

Brents’ Return

“For me to come back and actually step on a tennis court in the middle of July, then be declared fully recovered in August,” states Brents, “and then reach full speed on the platform tennis court in October was nothing short of miraculous.

“When I stepped onto the platform tennis court at the first tournament of the season in Philadelphia, my opponents looked at me as if they had seen a ghost,” exclaims Brents. “The opposition was totally unprepared for my return. They couldn’t gauge if I was there for serious competition or just for the fun of it.” The next day Brents and his partner reached the finals.

Brents, who is considered one of the quickest players in the game, even surprised his teammate. Partner Andy Kinney told him, “I watch you in the right-hand court, and can’t believe that you haven’t lost even a half a step.”

After the Rye tournament, the Brents-Kinney team was seeded number eight in the country. Brents would like to think his team has a shot at the championship. “Considering that I didn’t expect to be playing until January, our standing is terrific.”

Dr. Rozbruch and Mitzi Gerard offer the following tips to help prevent an Achilles tendon rupture:

• **10 to 15 minute warm-up of the calf and ankle muscles prior to play, using some type of aerobic activity:**

- running in place
- bicycling
- series of lateral, front, side-to-side movements

• **leg and calf muscle stretches**

- runner’s stretch
- stair stretch (heel extensions) - towel stretch

• **leg and calf muscle strengthening**

- resistive sports cord or rubber band exercises
- toe and heel raises.

Dr. Rozbruch also counsels that platform tennis players would do well to wear a slightly higher tennis shoe. “The higher the heel the less pull on the calf muscle during ambulation,” he explains.

For more on Dr. Rozbruch, visit www.JacobRozbruchMD.com